Acute Alcohol Intoxication Impairs Top–Down Regulation of Stroop Incongruity as Revealed by Blood Oxygen Level-Dependent Functional Magnetic Resonance Imaging

Ksenija Marinkovic,¹* Elizabeth Rickenbacher,² Sheeva Azma,³ and Elinor Artsy⁴

 ¹Department of Radiology, University of California, San Diego, California
 ²Champalimaud Foundation Neuroscience Programme at Instituto Gulbenkian de Ciência, Oeiras, Portugal
 ³Interdisciplinary Program in Neuroscience, Georgetown University, Washington, District of Columbia
 ⁴Postbaccalaureate Premedical Program, Tufts University, Medford, Massachusetts

Abstract: Functional neuroanatomy of executive functions has been delineated in a large number of neuroimaging studies using conflict-inducing tasks. The neural basis of alcohol's effects on cognitive control is poorly understood despite the evidence of impaired ability to evaluate competing demands and to inhibit maladaptive responses. To investigate the effects of moderate intoxication, healthy social drinkers participated in both alcohol (0.60 g/kg ethanol for men, 0.55 g/kg for women) and placebo conditions while being scanned using blood oxygen level-dependent (BOLD) functional magnetic resonance imaging (fMRI). A modified four-color Stroop task combined reading and color naming and used manual responses. Twenty subjects (10 women) were instructed to press a button corresponding to the font color except when a word was written in gray in which case they had to respond to the meaning of the word. Alcohol increased reaction times and a tendency to make more errors on incongruent trials. Behavioral indices of alcohol-induced premature responding correlated with the current drinking levels and impulsivity traits, suggesting an interaction between alcohol effects and personality predispositions. A distributed frontoparietal cortical network was activated by incongruity. However, moderate alcohol inebriation selectively attenuated anterior cingulate cortex (ACC) activation during both high-conflict trials and erroneous responses, indicating vulnerability of the regulative function subserved by the ACC. By disrupting top-down, strategic processing, alcohol may interfere with goal-directed behavior, resulting in poor self control. The present results support models proposing that alcohol-induced prefrontal impairments diminish inhibitory control and are modulated by dispositional risk factors and levels of alcohol consumption. Hum Brain Mapp 33:319–333, 2012. © 2011 Wiley Periodicals, Inc.

Key words: cognitive control; anterior cingulate; error-related activity

Contract grant sponsor: National Institutes of Health; Contract grant numbers: K01-AA13402, R01-AA016624, P41RR14075; Contract grant sponsor: Beverage Medical Research Foundation and Medical Investigation of Neurodevelopmental Disorders (MIND) Institute.

*Correspondence to: Ksenija Marinkovic, Department of Radiology, University of California, San Diego, 9500 Gilman Dr. 0841, La Jolla, California 92093-0841. E-mail: xenia@ucsd.edu Received for publication 16 March 2010; Revised 21 October 2010; Accepted 28 October 2010 DOI: 10.1002/hbm.21213

Published online 9 March 2011 in Wiley Online Library (wileyonlinelibrary.com).

© 2011 Wiley Periodicals, Inc.

INTRODUCTION

To successfully resolve certain conflict-inducing situations, it is necessary to inhibit responses that are normally automatic and habitual, but that cause interference under new constraints. This ability is termed "cognitive control" and is a hallmark of executive functions as we update and adjust to changing contingencies that indicate appropriate and safe behavior. One of the most popular tasks used to probe cognitive control is the Stroop Naming Task [Stroop, 1935] that manipulates interference by focusing on one feature of the stimuli while ignoring a dominant one. A series of color words are printed in different colors. On the Incongruent (INC) trials, the conflict arises when subjects are asked to focus on the font color while ignoring the habitual but irrelevant tendency to read the word. For instance, they are asked to respond "blue" when the word red is printed in blue. This conflict between word reading (READ; reflecting an automatic process) and color naming (reflecting a controlled process) results in increased error rates and slower reaction times [MacLeod, 1991].

This task has been used in a large number of imaging studies. Results of meta-analyses of activation patterns evoked by the Stroop task suggest that conflict-induced functional activations are subserved by a predominantly frontal cortical network. Main points of convergence across a large number of studies are bilateral dorsal anterior cingulate cortex (ACC), bilateral but left-dominant inferior frontal junction (IFJ), inferior frontal gyrus (IFG), inferior parietal lobule (IPL), and presupplementary motor area (pre-SMA) [Laird et al., 2005; Neumann et al., 2005]. This activation pattern may not be specific to the Stroop task, because a highly overlapping prefrontal network is activated by a range of cognitive tasks [Duncan and Owen, 2000]. Nevertheless, ACC, also termed rostral cingulate zone [Ridderinkhof et al., 2004], holds a central position in that network. It is activated in a variety of tasks as a function of task difficulty, response conflict, novelty, and anticipation [Bush et al., 2000; Paus, 2001]. As it is particularly sensitive to tasks in which automatic responses need to be overcome in favor of task-relevant, but nonautomatic responses, the ACC seems to underlie controlled processing in a sense of guiding behavior that is not well rehearsed or habitual [Procyk et al., 2000; Raichle et al., 1994]. This multifaceted top-down role of the ACC in goaloriented actions is supported by its anatomical connections with dorsolateral prefrontal cortex (dlPFC), motor cortex, spinal cord, and limbic structures [Barbas, 2000; Bush et al., 2000; Devinsky et al., 1995]. Furthermore, functional magnetic resonance imaging (fMRI) studies indicate its importance in error monitoring and behavioral adjustments [Carter et al., 1998; Ullsperger and von Cramon, 2004]. The temporal dimension of error-related responses has been investigated with event-related potentials. The error-related negativity (ERN) component, presumably generated in the ACC, is evoked on erroneous trials [Coles et al., 1995; Holroyd and Coles, 2002]. This deflection is attenuated by

alcohol [Easdon et al., 2005; Holroyd and Yeung, 2003; Ridderinkhof et al., 2002], suggesting vulnerability of the error-monitoring network to alcohol intoxication.

Neuroimaging data delineating alcohol effects on cognitive control are scant in spite of the crucial importance of understanding the neural basis of alcohol's effects on behavioral self-regulation. Gundersen and colleagues [2008] observed decreased activation in ACC and cerebellum during a working memory task under a moderate alcohol dose. A slightly lower alcohol dose and a different variant of a working memory task resulted in increased activation in dlPFC and decreased activation in parietal regions [Paulus et al., 2006]. Impairments in a distributed, but primarily frontal circuitry were observed under conditions of simulated driving while intoxicated [Calhoun et al., 2004; Meda et al., 2009]. Using Stroop task and event-related potentials, Curtin and Fairchild [2003] reported reduced frontal components to incongruity under alcohol, suggesting impairment in evaluative and regulative processes.

Neuroimaging studies of abstinent chronic alcoholics have observed decreased frontal activation during cognitive control tasks. In a PET study, mediofrontal hypometabolism correlated with response speed on Stroop task in abstinent alcoholics [Dao-Castellana et al., 1998]. Hypoactivity of dIPFC areas was also observed in a cohort of chronic alcoholics with fMRI during a stop-signal task, suggesting impaired impulse control [Li et al., 2009]. These results are in agreement with other lines of research, suggesting that chronic alcoholism results in deficits in executive functions [Oscar-Berman and Marinkovic, 2007], with prefrontal cortex being particularly vulnerable to alcoholinduced damage [Sullivan and Pfefferbaum, 2005]. It has been proposed that the prefrontal deficits underlie inability to control drinking [Kalivas, 2009], which is one of the criteria for alcohol dependence (DSM-IV; [APA, 1994]).

It is generally accepted that a decreased ability to inhibit prepotent responses in favor of the rewarded ones relates to the inability to refrain from drinking and may be a factor in subsequent alcohol or drug abuse [Fillmore, 2003; Finn, 2000; Jentsch and Taylor, 1999; Lyvers, 2000]. Moreover, impaired cognitive control is influenced by genetics, and it contributes to the vulnerability to alcoholism [Schuckit, 2009]. In spite of their relevance, the effects of acute intoxication on the neural basis of cognitive control are poorly understood. We used fMRI to investigate alcohol effects on cognitive control as reflected in the ability to inhibit prepotent responses in favor of the rewarded ones. We used a modified version of the Stroop task that combined color-naming and READ and that contained trials that were high in conflict (i.e., incongruous), low in conflict (i.e., congruous), or neutral (NEUT, i.e., noncolor words). Healthy, young social drinkers served as their own controls by participating in both moderate alcohol and placebo conditions in a counterbalanced manner. We investigated beverage-related impairments in patterns of behavioral and neural activity as a function of taskinduced conflict. In addition, we examined effects of

inebriation on error-related activity with particular emphasis on the anterior cingulate contributions, given previous reports of attenuated ERN under alcohol [Ridderinkhof et al., 2002] and the assumed ACC generators [Ullsperger and von Cramon, 2001]. Taking advantage of the excellent spatial resolution of the fMRI method, we tested the hypothesis of an alcohol-specific decrease in ACC activity on high conflict and erroneous trials. Gender is an important factor mediating alcohol's effects on cognitive control [Fillmore and Weafer, 2004]. Higher prevalence of alcohol-use disorders among men may result from a confluence of genetic, biological, and social factors [Nolen-Hoeksema and Hilt, 2006], calling for more research to delineate their respective influences. Consequently, potential gender differences in alcohol's effects on the behavioral and physiological variables were explored in the current study.

METHODS

Research Participants

Twenty individuals (10 women) participated in both alcohol and placebo sessions in a counterbalanced manner. Average age (\pm st. dev.) was 24.9 \pm 3.6 years and age range was 21-35 years. They were all right-handed, nonsmoking native English speakers with no alcohol- or drugrelated problems. None of the participants had any healthrelated problems and none were taking any medications. The participants were predominantly white, but the sample included one Pacific Islander and two Hispanic individuals. On average, participants reported light-to-moderate drinking pattern imbibing 1.8 ± 1.2 times per week on average, 2.5 ± 1.0 drinks per occasion. Men and women did not differ in the amount or frequency of drinking. No alcoholismrelated symptoms were detected with the Short Michigan Alcoholism Screening Test [Selzer et al., 1975], and the subjects reported no family history of alcoholism or drug abuse (first and second degree relatives). Their responses on personality questionnaires were in the normal range.

Task

A modified Stroop paradigm combined trials of READ and color naming in a randomized manner. It consisted of four conditions that varied interference between font color and word meaning (see Fig. 1). Responses were given manually on a four-button response box. Response conflict was elicited by the INC condition (e.g., word "red" printed in blue color), in comparison with *congruent* (CONG) trials on which the meaning corresponds to the display (e.g., word "red" printed in red color). The NEUT condition consisted of common noncolor words (e.g., word "paid" written in color) that were matched with the color words in length and lexical frequency [Francis and Kucera, 1982]. Words in the INC, CONG, and NEUT conditions were written equiprobably in four colors (red, green, yel-





Trial examples of the four conditions in this modified Stroop paradigm, along with the correct response color for each trial. The task included the Congruent (CONG) condition in which the font color coincides with the word meaning, INC condition that elicits response conflict due to incongruity between the font color and the word meaning, and Neutral (NEUT) condition in which common words were written in color. Participants were asked to press a button corresponding to the color of the font whenever a word was written in color. Words in the Read condition were written in gray, and participants had to press a button corresponding. Trials were presented for 300 ms every 2 s in a randomized manner.

low, and blue) and represented 55.5% (18.5% each) of all the trials. Subjects had to respond to the font color of these words. An additional READ control condition (44.5% trials) consisted of color words written in gray to which subjects pressed the button corresponding to the meaning of the word. The purpose of this condition was to maintain READ dominance and automaticity [Repovs, 2004]. Subjects were instructed to press a button corresponding to the font color whenever they saw a word written in color and to press a button corresponding to the word meaning when it was written in gray. They were asked to respond on each trial as quickly as possible without losing accuracy while using index and middle fingers of both hands on a four-button magnet-compatible response box. They practiced response mapping with a series of progressively realistic practice runs during a familiarization session. At first, they practiced responding to strings of x's written in each color, followed by responding to words written in gray. Finally, they were given a practice run resembling the real experiment and a test run to ascertain that their performance was at perfect or near-perfect levels, that is, at or above 95% accuracy. Performance accuracy and speed were analyzed with a mixed model ANOVA with gender as a between-group factor and beverage and conditions as within-subject factors [Woodward et al., 1990; Fig. 2]. In addition to the F-values and probability, Cohen's d statistic [Cohen, 1988] corrected for sample size bias resulting in Hedges' d [Hedges and Olkin, 1985] is reported for all mean value comparisons in the text and Tables I and II.

The words were presented for 300 ms on the black background followed by a fixation string (a series of x's) with a total trial length of 2 s. The stimuli were shown in the center of a rear-projection presentation screen in a manner synchronized with the scanner using the Presentation



Stroop color-word interference effect is reflected in longer RTs and lower accuracy on INC trials when compared with other conditions overall. When intoxicated, participants had longer average RTs and tended to make more errors on INC trials. Alcohol versus placebo comparisons: *P < 0.05, \$P < 0.07.

software package (Neurobehavioral Systems). Each subject was presented with five 5-min runs that composed a total of 100 INC, 100 CONG, 100 NEUT, and 240 READ trials. In addition, 190 fixation trials were randomly intermixed providing the temporal jitter needed for optimal deconvolution of the BOLD signal [Burock et al., 1998]. Optimized randomization of the event-related sequence was achieved with the Optseq program within the FS-FAST software (http://surfer.nmr.mgh.harvard.edu/optseq).

Experimental Design and Procedure

Subjects served as their own controls by participating in both alcohol and placebo sessions in a counterbalanced manner. During an introductory session, subjects were familiarized with the laboratory setting, experimental procedure, and the task, minimizing the potential effects of situation-induced arousal. Subjects also provided more detailed information about their medical history, family history of alcoholism, level of response to alcohol [Schuckit et al., 1997], severity of their alcoholism-related symptoms [Selzer, 1971], quantity and frequency of alcohol use [Cahalan et al., 1969], and handedness [Oldfield, 1971]. To obtain a comprehensive dispositional profile for each subject particularly with respect to disinhibitory, novelty seeking, and socialization traits, the following questionnaires were used: Childhood Hyperactivity Questionnaire, [Tarter et al., 1977]; Eysenck Personality Questionnaire, EPQ [Eysenck and Eysenck, 1975]; Socialization Scale of the California Psychological Inventory [Gough, 1994]; Zuckerman Sensation Seeking Scale [Zuckerman, 1971]; Evsenck Impulsiveness and Venturesomeness Scale [Eysenck and Eysenck, 1978]. Correlations between scores on personality questionnaires and performance measures were calculated as a function of beverage and described below.

Written informed consent approved by the Human Research Committee at Massachusetts General Hospital and the Partners Healthcare Network was obtained from all subjects before participation. Subsequently, the subjects

participated in placebo and alcohol sessions that were counterbalanced in order of presentation. Alcohol was given in the first session to 10 of 20 subjects. The two sessions were 30 \pm 26 days apart on average. Upon their arrival to the laboratory, the subjects were queried about their compliance with the requirement to abstain from food for 3 h and from alcohol at least 48 h before each experimental session. Female subjects were given a pregnancy test each time to ascertain that they were not pregnant. Blood alcohol concentration (BAC) was measured with a breathalyzer (Draeger) upon arrival and throughout the session when the subjects were outside the scanner. During the actual scan, we used Q.E.D. Saliva Alcohol Test (OraSure Techn) to estimate the BAC, because no electronic device can be used in the scanner chamber. Identical procedure was used in both alcohol and placebo sessions with the exception of the beverage contents. The subjects rated their moods and feelings with the adapted Biphasic Alcohol Effects Scale [BAES, Martin et al., 1993] on three occasions: before drinking, on the ascending BAC limb (immediately prior to task performance) and on the descending limb (immediately after the scan). They were asked to rate their momentary feelings on a series of adjectives probing stimulant (e.g., vigorous) and sedative (e.g., sluggish) alcohol effects as well as how tired, worn-out, high, euphoric, or sexy they felt at the time. The scores on BAES scale were analyzed with a mixed model ANOVA with gender as a between-group factor and beverage (alcohol and placebo) and phase (before experiment, ascending BAC, and descending BAC) as within-subject factors on each subscale separately.

In each session, either alcohol [0.60 g/kg for men, 0.55 g/kg for women, presented as cocktail containing vodka (Grey Goose, Bacardi) as 20% v/v in orange juice] or placebo (the same volume of orange juice) was administered within a single-blind procedure [Marinkovic et al., 2004] and consumed in about 10 min. Gender-specific dosing was used in order to adjust for the body mass index difference [Breslin et al., 1997]. On average, participants consumed ~2.8 standard drinks defined as 1.5 fl oz of vodka.

TABLE I. Summary of beverage effects on behavioral
measures as reflected in group means ± standard
deviations for placebo and alcohol conditions, F- and
P-values of the difference and Hedges' unbiased d
effect size index

	Placebo	Alcohol	$F_{(1,18)}$	<i>P</i> -value	D-value					
Performance										
Accuracy										
Inc	92.2 ± 4.9	89.8 ± 5.4	3.8	< 0.07	0.46					
Cong	96.3 ± 2.8	94.6 ± 5.2	1.6	< 0.22	0.39					
Neut	95.6 ± 3.4	94.3 ± 6.3	0.8	< 0.39	0.25					
Read	95.6 ± 2.5	94.2 ± 4.8	2	< 0.17	0.36					
All cond	94.9 ± 3.8	93.2 ± 5.7	2.52	< 0.13	0.35					
RTs										
Inc	828.6 ± 86.8	851.3 ± 97.5	3.6	< 0.07	0.24					
Cong	682.3 ± 80.6	705.7 ± 80.1	5.4	< 0.05	0.29					
Neut	721.2 ± 72.1	743.9 ± 71.8	4.5	< 0.05	0.31					
Read	726.8 ± 85.3	749.7 ± 88.9	3.8	< 0.07	0.26					
All cond	739.7 ± 96.7	762.6 ± 99.6	5.6	< 0.05	0.23					
BAES scales										
Stimulation										
Baseline	24.8 ± 13.6	26.3 ± 13.6	0.24	>0.5	0.11					
Asc BAC	23.8 ± 14.3	28.5 ± 13.8	3.5	< 0.08	0.33					
Desc BAC	17.8 ± 12.1	16.1 ± 13.8	1	< 0.3	0.13					
Sedation										
Baseline	15.5 ± 12.1	15.0 ± 9.8	0.1	>0.5	0.05					
Asc BAC	14.8 ± 11.3	17.6 ± 11.9	1.7	< 0.2	0.23					
Desc BAC	22.9 ± 14.7	31.5 ± 12.9	16.6	< 0.001	0.59					
High										
Asc BAC	1.5 ± 2.1	3.2 ± 2.7	12	< 0.01	0.66					
Euphoric										
Asc BAC	2.1 ± 2.1	3.05 ± 2.9	6.2	< 0.05	0.37					
Sexy										
Asc BAC	2.3 ± 2.5	2.7 ± 3.0	3.8	< 0.07	0.15					
Tired										
Desc BAC	4.7 ± 2.5	5.7 ± 2.7	9.1	< 0.01	0.39					
Post experimental scales										
Likert 1–5										
Task diff.	3 ± 0.8	2.9 ± 0.8	0.14	>0.5	0.06					
Nausea	1.2 ± 0.5	1.2 ± 0.4	0.0	>0.5	0.00					
Dizziness	1.2 ± 0.4	1.9 ± 1.1	11.4	< 0.01	0.81					
Bev. content	1.4 ± 0.9	4.5 ± 0.9	57.1	< 0.0001	1.66					
Intoxication	1.1 ± 0.2	2.8 ± 0.9	70.2	< 0.0001	1.57					
Scale 0–5										
No. of drinks	0.1 ± 0.4	2.3 ± 0.9	64.2	< 0.0001	1.63					
BAC										
Before task	0.0 ± 0.0	0.043 ± 0.01	n/a	n/a	n/a					
After task	0.0 ± 0.0	0.052 ± 0.01	n/a	n/a	n/a					

Included are measures of performance (accuracy and RTs) for each condition separately as well as average across conditions (All cond); BAES scales (stimulant, sedative effects, and self-reports on high, euphoric, sexy, and tired scales for the baseline (before experiment), ascending and descending BAC; postexperimental self-reports on Likert scales (1–5) for task difficulty, nausea, dizziness, beverage content, and intoxication. Participants also estimated how many alcoholic drinks were contained in the beverage on a scale ranging from 0 to drinks with 0.5 drinks increments. Finally, BAC measures are also included. The task was administered 44 \pm 9 min after the subjects were presented with their drink. The average BAC measured before the task was 0.043% \pm 0.01% and reached 0.052% \pm 0.01% upon its completion, indicating that the task was administered on the ascending BAC limb. It was followed by another task. At the end of each experimental session, the participants filled out a detailed questionnaire about perceived task difficulty, type, and content of the beverage they imbibed, about how intoxicated, nauseous, or dizzy they felt. Transportation home was provided to all participants upon completion of each session.

Image Acquisition and Analysis

Imaging data were acquired with a 3 T Siemens Trio whole-body scanner (Siemens, Erlangen, Germany). Special care was taken to minimize head motion with the use of a special pillow, foam padding, and head "clamps" that also allowed subjects to maintain a comfortable position during scanning. Exposure to scanner noise was reduced with 29-db earplugs and pillow padding. Subjects could view the stimuli comfortably as they were rear-projected onto a mirror fitted onto the head coil. For each subject, two high-resolution 3D MPRAGE (magnetizationprepared rapid gradient echo) T1-weighted sequences that optimize contrast for a range of tissue properties were obtained with the following parameters: TR = 2.53 s, TE = 3.25 ms, flip angle = 7° , FOV = 256, 128 sagittal slices, 1.33 mm thickness, and in-plane resolution 1×1 mm. These two high-resolution structural images were used for spatial normalization and cortical surface reconstruction. A series of functional whole-brain BOLD images was collected using a T2*-weighted EPI sequence of 28 interleaved 5-mm thick slices in axial-oblique AC-PC orientation with TR = 2 s, TE = 30 ms, flip angle = 90° , FOV = 200 mm, matrix = 64 \times 64, resulting in 3.13 \times 3.13 inplane resolution.

The FreeSurfer and FS-FAST (Free-Surfer-Functional Analysis Stream) analysis package was used to analyze brain images [Burock and Dale, 2000; Dale et al., 1999; Fischl et al., 1999a]. Each subject's cortical surface was reconstructed using an automatic gray/white segmentation, tessellation, and inflation of the folded surface patterns (http://surfer.nmr.mgh.harvard.edu/). These surfaces were registered with a canonical brain surface created from an average of 40 brains [Fischl et al., 1999b], allowing for high-resolution group averaging based on surface alignment. Functional data were motion corrected with AFNI software [Cox, 1996; Cox and Jesmanowicz, 1999], spatially smoothed with a 3D 5-mm Gaussian fullwidth half-maximum filter, corrected for temporal drift, and normalized to correct for signal intensity changes. The data were carefully checked for motion or artifacts. Motion for one subject's run in one session was 3.25 mm and was excluded, along with the matching run in the complementary session for the same subject. Otherwise, the motion

Area	Talairach coord	Condit. (F _{3,51})	INC–Alc, $F_{1,17}$	d	INC–Plac, $F_{1,17}$	d
Left IFG	-42.9 20.7 17.0	26.9 < 0.0001	16.4 < 0.001	0.97	34.1 < 0.0001	1.08
Left IFJ	-52.1 3.5 28.1	18.7 < 0.0001	15.3 < 0.001	0.78	30.7 < 0.0001	0.97
Left IPC	-29.3 -45.9 45.1	16.4 < 0.0001	23.6 < 0.0001	0.69	23.5 < 0.0001	1.10
Left sup PF	$-20.9\ 1.5\ 44.2$	12.1 < 0.0001	21.6 < 0.001	0.39	19 < 0.001	0.74
Left aINS	-9.9 4.7 45.1	24.2 < 0.0001	32.5 < 0.0001	0.99	47.4 < 0.0001	1.06
Left ACC	-6.2 24.1 21.8	3.8 < 0.01	Ns	0.16	$14.9 < 0.01^*$	0.72
Left SMA	-9.0 12.6 42.5	11.8 < 0.0001	7.4 < 0.01	0.42	28.3 < 0.0001	0.54
Right IFJ	58.6 5.5 20.5	7.8 < 0.001	8.5 < 0.01	0.35	21.5 < 0.001	0.71
Right IPC	27.8 -47.1 43.3	16.1 < 0.0001	9.5 < 0.01	0.41	21.4 < 0.001	0.76
Right aINS	31.7 18.6 -1.7	10.1 < 0.0001	10.8 < 0.01	0.59	31.3 < 0.0001	0.78
Right ACC	9.3 30.3 18.6	4.7 < 0.01	6.7 < 0.01	0.35	$8.1 < 0.01^*$	0.58
Right SMA	8.9 25.5 39.6	5.5 < 0.01	8.7 < 0.01	0.13	17.4 < 0.001	0.47

 TABLE II. Summary of statistical results for ROIs including their Talairach coordinates, significance of the main effect of Condition, and comparisons of INC vs all other conditions for alcohol and placebo

Included are F- and P-values of the difference and Hedges' unbiased d effect size index.

*The only ROIs with significant alcohol-induced attenuation of the activity to INC were in the left ACC ($F_{1,17} = 4.7$, P < 0.05, d = 0.63) and right ACC ($F_{1,17} = 8.2$, P < 0.01, d = 0.56).

IFG, inferior frontal gyrus; IFJ, inferior frontal junction; IPS, intraparietal sulcus; sup PF, superior prefrontal cortex; aINS, anterior insula; ACC, anterior cingulate cortex; SMA, supplementary motor area.

did not exceed 2 mm for any subject or run and did not differ between sessions. Event-related hemodynamic responses (HDRs) were estimated with a finite impulseresponse model (FIR) [Burock and Dale, 2000], which provides unbiased estimates of the average signal intensity at each time point for each trial type and makes no a priori assumptions about the shape of the HDR. To capture the timecourse of the BOLD activation, the mean and variance values of the HDR were estimated for each TR within a time window of 18 s. Motion parameters derived from realignment correction were entered into the model as regressors. F-distributed statistical activation maps were generated from averaged responses for each contrast of interest and resampled onto the common cortical surface space for the surface-based analysis, which included trials with correct responses. Contrasts were generated for each condition versus fixation in order to assess the overall activity, whereas conflict-specific activity was examined by contrasting INC verus other conditions. Group-average results were obtained using a random-effects statistical model, which takes into account the intersubject variance, allowing for inferences to the population [Friston et al., 1999], and projected onto an inflated brain with average curvature [Fischl et al., 1999b]. Potential baseline shifts were removed by subtracting the average hemodynamic response before stimulus onset from the hemodynamic response waveform for both "placebo" and "alcohol" conditions, equating their respective baselines. The resulting statistical parametric maps of the voxel-wise group-average analysis are presented in Figures 3 (for the overall activity) and 4 (for the conflict-specific activity). In addition, region-of-interest (ROI) analyses were conducted in order to further explore the main effects and interactions of the factors of gender, beverage, and condition. The ROIs were based on functional features and defined on a groupaverage voxel-wise analysis using the unbiased orthogonal contrast (i.e., all contrasts vs. fixation) [Friston et al., 2006; Poldrack, 2007]. ROIs included voxels active at P < 0.0001at the activity peak (TR window 4-6 s latency). The ROI placement was blind to each subject's activation patterns, because the group-average-based ROIs were automatically transferred from the average cortical surface onto each individual's surface. Percent signal change from baseline was computed for each ROI and each subject, session, and condition and presented in the form of timecourses that were also baseline normalized. The ROIs and groupaverage timecourses for alcohol and placebo are presented in Figure 5. ANOVA analyses were conducted on these values using a mixed model with a between-group factor of gender and within-subject factors of beverage and condition. The reported significance values have been modified to reflect Huynh-Feldt adjustment for the Condition factor, because it contains four levels. One male subject's data were unusable in one condition, because he fell asleep in the scanner, and so the neuroimaging results are reported for N = 19 (10 females).

Error Analysis

To explore error-related activity, trials with erroneous responses were matched with correct trials that belonged to the same stimulus type. Half of the matched correct trials preceded and half followed error trials but were never ordinally adjacent as they were separated by intervening trials. The FIR estimates were obtained as described earlier, and group averages were calculated with randomeffects statistical model. Results of the voxel-wise analysis at 4–6 s latency are shown in Figure 6, left panel, for the medial surface of the left and right hemispheres. As



Figure 3.

Voxel-wise group-average statistical maps are displayed on the inflated lateral and medial cortical surfaces for both hemispheres and for all conditions under placebo and alcohol at the activity peak (4–6 s latency). The ACC activity evoked under placebo was significantly attenuated by alcohol (black arrows). The color bar denotes *P*-values obtained with the random effects group analysis of all conditions versus fixation contrasts.

described earlier, ROIs were obtained based on the groupaverage data and automatically transferred on each subject's cortical surface. Baseline-normalized % signal change values for the factors of error (error vs. correct) and beverage (alcohol vs. placebo) were entered into a withinsubject ANOVA (Fig. 6, right panel).

Nine subjects (3 males) that were included in the analysis made 26.2 \pm 17.1 errors on average, with the minimum



Figure 4.

Voxel-wise group-average maps obtained with the random effects group analysis for INC versus all other conditions contrasts and displayed on the inflated lateral and medial cortical surfaces for both hemispheres and beverage conditions at 4–6 s latency. Conflict-evoked activity observed in the ACC was stronger under placebo (black arrows) than alcohol.



Figure 5.

Shown are group-average timecourses (% signal change) of the BOLD activation for all conditions and both beverages for a set of ROIs. The ROIs (delineated in white) were derived from group-average voxel-wise analysis using the unbiased orthogonal contrast and were baseline normalized. Conflict-related activity was attenuated significantly in the ACC. IFJ: inferior frontal junction; IFG: inferior frontal gyrus; IPS: intraparietal sulcus; SMA: supplementary motor area; ACC: anterior cingulate cortex.

of 15. Possible differences between the participants who did or did not have sufficient number of errors were carefully examined on the scales of task difficulty, intoxication, moods, personality, and BAC. Subjects who were included in the "error" analysis rated the task as being more difficult under intoxication on the Likert 1–5 scale (3.4 \pm 0.6) than the participants who made fewer errors (2.5 \pm 0.7), $F_{(1.18)} =$ 12.6, P < 0.01, d = 1.27. On the BAES scale, the participants generating more errors reported being less stimulated on the ascending BAC limb (21.9 \pm 14.1) than the individuals with fewer errors (33.9 \pm 11.5), $F_{(1,18)} = 4.4$, P < 0.05, d =0.83. No other differences were detected as the groups did not differ in terms of self-reported intoxication, sedation, BAC, drinking habits, or on any personality measures. These results suggest that increased error rates are not due to inherent personality traits, but may result from situational variations in task difficulty and alcohol-induced moods.

Posterror behavioral adjustment is commonly reflected in slower RTs following incorrect responses [Jentzsch and Dudschig, 2009; Rabbitt, 2002]. To examine this phenomenon, the RT analysis included all error trials that were closely followed by a correct response that was strictly matched for the condition type. The errors and posterror correct responses were 4.9 ± 1.1 s apart. The posterror correct responses were further matched with correct responses belonging to the same condition that were close to the error trial but were, on average, 13.0 ± 2.2 s apart. Reaction times were analyzed with repeated measures ANOVA including within-subject factors of beverage (alcohol and placebo) and trial type (errors, posterror correct, and other correct trials).

RESULTS

Behavioral Measures

Performance

Our task successfully manipulated the Stroop interference effect as indicated by lower accuracy ($F_{1,18} = 24.5$,



Figure 6.

Group-average voxel-wise random effects analysis of erroneous and correct trials versus fixation. Significant activity was observed only on medial views of the inflated cortical surfaces as shown in the left figure panel. Right panel: average % signal change (\pm s.e.m.) from baseline for the bilateral ROIs centered on the ACC, showing that error-induced activity is significantly attenuated under intoxication.

P < 0.0001, d = 0.82) and slower RTs ($F_{1,18} = 205.5, P < 0.0001$ 0.0001, d = 1.18) on INC trials when compared with all other conditions (see Fig. 2). Table I lists behavioral results for all beverage comparisons. With the exception of a tendency to make more errors on INC trials under alcohol compared to placebo ($F_{1,18} = 3.8$, P < 0.07, d = 0.46), there were no other beverage or gender effects on performance accuracy. In contrast, RTs were significantly longer under alcohol (762.6 \pm 99.6 ms) than under placebo (739.7 \pm 96.7 ms) as indicated by the main effect of beverage $(F_{1,18} = 5.6, P < 0.05, d = 0.23)$. Furthermore, the main effect of condition ($F_{3,54} = 125$, P < 0.0001) was due to the overall fastest responses on CONG trials, 694.0 \pm 80.2 ms $(F_{1.18} = 208, P < 0.0001, d = 0.76)$, and slowest responses on INC trials, 840.0 \pm 91.8 ms ($F_{1,18} = 205.5$, P < 0.0001, d = 1.18), when compared with other conditions. RTs on NEUT and READ trials did not differ (732.5 \pm 71.9 ms; 738.2 \pm 86.7 ms, respectively) but were longer than the congruous ($F_{1,18} = 64.2$, P < 0.0001, d = 0.50) and shorter than the incongruous ($F_{1,18} = 148.8, P < 0.0001, d = 1.06$) across both beverage conditions.

Mood ratings

There were no gender effects on any of the measures. Overall, subjects felt more sedated and less stimulated at the end of the scan, as indicated by the main effect of phase for both stimulation ($F_{2,36} = 22.3$, P < 0.0001) and sedation ($F_{2,36} = 22.5$, P < 0.0001) subscales. Subjects tended to rate themselves as more stimulated on the ascending BAC under intoxication ($F_{1,18} = 3.5$, P < 0.08, d = 0.33) and were more sedated on the descending BAC ($F_{1,18} = 16.6$, P < 0.001, d = 0.59) when compared with placebo. This was confirmed with additional mood probes as subjects reported to be significantly more high ($F_{1,18} = 16.6$).

12.0, P < 0.01, d = 0.66) and euphoric ($F_{1,18} = 6.2$, P < 0.05, d = 0.37) and tended to feel more sexy ($F_{1,18} = 3.8$, P < 0.07, d = 0.15) on the ascending BAC when given alcohol, when compared with placebo. Conversely, they felt significantly more tired ($F_{1,18} = 9.1$, P < 0.01, d = 0.39) on the descending BAC when comparing alcoholic versus placebo beverage conditions.

Postexperimental questionnaire

Subjects provided ratings on Likert scales (1-5). Each scale was analyzed with a mixed model ANOVA with gender as a between-group and beverage as a within-subject factor. Participants rated the task as being moderately difficult (2.9 \pm 0.7), but the perceived difficulty was not affected by the beverage. On the scale from 1 (definitely contains no alcohol) to 5 (definitely contains alcohol), participants rated the beverage contents as 4.5 \pm 0.9 under alcohol and 1.4 ± 0.9 under placebo, and these ratings differed significantly ($F_{1,18} = 57.0$, P < 0.0001, d = 1.66). They estimated that the alcoholic beverage contained 2.3 \pm 0.9 "alcoholic drinks," which was a slight underestimate of the actual amount containing 2.8 standard drinks on average defined as 1.5 fl oz of vodka. They also reported that the placebo beverage contained 0.1 \pm 0.4 "alcoholic drinks." No gender effects were found on any of these measures. However, there was a significant interaction between the factors of gender and beverage ($F_{1,18} = 4.6$, P < 0.05) for the intoxication self-reports. Based on 1–5 Likert scale, women felt more intoxicated than men when given alcohol ($F_{1,18} = 4.8$, P < 0.05, d = 0.72), as they reported being "moderately intoxicated" (3.2 \pm 0.8), whereas men were only "slightly intoxicated" (2.4 \pm 0.8). Even though being intoxicated did not affect the nausea ratings on the scale of 1 (not at all) to 5 (very much; overall rating was 1.2 \pm 0.4), subjects reported feeling slightly but significantly dizzier under alcohol (1.9 \pm 1.1) when compared with placebo conditions (1.2 \pm 0.4; $F_{1,18} =$ 11.4, P < 0.01, d = 0.81).

Personality questionnaires

Beverage-related differences in the number of self-corrections (trials on which participants corrected themselves) correlated with two measures of impulsivity: Psychoticism Scale of Eysenck's Personality Questionnaire (r = 0.76, P < 0.001) and Impulsivity Scale on Eysenck's Impulsiveness and Venturesomeness Scale (r = 0.54, P < 0.01), possibly suggesting that, as the baseline impulsivity increases, alcohol causes more impairment in the ability to inhibit the prepotent but erroneous responses.

Neuroimaging Results

Task-induced activity

Voxel-wise analysis was performed using random effects model of the group data. Figure 3 shows the overall activity to all four stimulus types during placebo and alcohol conditions. Conflict-evoked activity (i.e., INC vs. other stimulus types) is presented in Figure 4, indicating a subset of areas that contribute to interference-related activity. For the ROIs representing main foci of cortical activity, additional statistical analyses were performed on activity levels as reflected in baseline-normalized percent signal change values. A distributed cortical network was activated by this interference task, in agreement with results from other fMRI studies using Stroop paradigm summarized with meta analyses [Laird et al., 2005; Neumann et al., 2005]. Significant effects are listed in Table II, and a subset of timecourses is shown in Figure 5.

The most prominent effect of alcohol on baseline-normalized percent signal change measured in ROIs was attenuation of conflict-related activity on the INC trials in ACC regions (Table II). Activity to INC under placebo was significantly stronger than activity under alcohol both in the left ($F_{1,17} = 4.7, P < 0.05, d = 0.63$) and right ACC $(F_{1.17} = 8.2, P < 0.01, d = 0.56)$. The main effect of beverage was statistically significant in the right ACC ($F_{1,17}$ = 4.6, P < 0.05), with stronger activity under placebo than alcohol (0.11 vs. 0.08% signal change). This alcoholinduced attenuation of activity on conflict trials was limited to ACC bilaterally. The main effect of Condition was observed in distributed, but primarily frontoparietal areas with significantly stronger activity to INC than to other stimulus conditions (Table II). CONG, NEUT, and READ did not differ in evoked signal strength with one exception. Activity in the left IFG showed graded responses to the four conditions, so that the activity to INC was larger than to NEUT ($F_{1,17} = 9.9$, P < 0.01, d = 0.68), which, in turn, was larger than activity to CONG ($F_{1,17} = 14.1$, P < 14.1 0.01, d = 0.54). Activity to CONG was only marginally stronger than the activity to READ ($F_{1,17} = 2.0$, P < 0.06, d = 0.25; Fig. 5). As discussed below, this unique pattern may be due to the lexicosemantic function ascribed to this area [Wagner et al., 2001]. Effects of gender were limited to the left ACC where men showed stronger overall activity than women ($F_{1,17} = 10.5$, P < 0.01, d = 0.91).

Error-related activity

Repeated measures ANOVA was carried out on RTs as a function of beverage and correct versus error responses. Average RTs of the trials included in the imaging analysis were not significantly affected by beverage and did not differ between the errors (798.6 \pm 91.9) and correct responses (804.1 \pm 96.9).

To test the hypothesis of alcohol effects on the neural basis of error processing, the activity generated on error and matched correct trials was analyzed in ACC bilaterally. Figure 6 shows voxel-wise analysis for both alcohol and placebo conditions as well as average % signal change differences for the ROIs in ACC. The ROI analysis (Fig. 6, right panel) indicates that the ACC activity is significantly stronger on error when compared with correct trials, but only under placebo ($F_{1,8} = 6.0$, P < 0.05, d = 0.72) and not alcohol ($F_{1,8} = 0.2$, P > 0.5, d = 0.18). Thus, moderate alcohol inebriation attenuates both conflict-related and error-related activity in ACC area.

Comparison of RTs on the subset of correct trials that closely followed erroneous responses and were matched by condition type with RTs on other matched correct trials and errors indeed confirmed posterror slowing. Beverage did not affect RTs significantly on the three trial types (i.e., error, posterror correct, and other correct). Erroneous RTs (829.4 \pm 168.7) did not differ from either posterror corrects or other correct responses, whereas the posterror correct responses (819.3 \pm 120.0) were indeed significantly longer than other correct responses (760.0 \pm 77.2), $F_{(1,8)} = 6.4$, P < 0.05, d = 0.56, confirming other reports of posterror slowing [Jentzsch and Dudschig, 2009; Rabbitt, 2002].

DISCUSSION

Our study indicates that a distributed, frontoparietal cortical network is activated by incongruity during the Stroop task, confirming other reports [Laird et al., 2005; Neumann et al., 2005]. More importantly, however, activity in the anterior midcingulate area is selectively attenuated by moderate alcohol inebriation during both high-conflict trials and erroneous responses. This finding indicates that the regulative, top–down function subserved by the ACC is vulnerable to moderate intoxication [Ridder-inkhof et al., 2002]. The importance of the ACC for cognitive control has been emphasized in a large number of studies showing that it contributes to better performance by maintaining decision contingency in the active

attentional focus, by inhibiting inappropriate response tendencies, or by performance monitoring [Bush et al., 2000; Rushworth et al., 2004]. The ACC may subserve the topdown regulation necessary to maintain performance in high-conflict situations in which prepotent responses need to be overcome in favor of nonautomatic, but task-relevant responses [Procyk et al., 2000; Roelofs et al., 2006]. An appropriate level of cognitive control ensures that our actions are in agreement with our intents and goals, and that they fit with in the context. It also includes suitable planning and self-regulation, which are necessary to avoid succumbing to distractions, and permits refraining from temptations such as heavy drinking. In contrast to automatic processing that is strongly influenced by the external stimuli, controlled processing comprises effort to overcome habitual responses and perform less practiced tasks [Schneider and Chein, 2003]. Alcohol disrupts the topdown, strategic mode of processing that is regulated by prior goals and intentions and makes us more susceptible to the momentary and immediate cues. Indeed, it is generally accepted that impaired self-regulation of executive functions such as an inability to refrain from drinking may be a factor in subsequent alcohol or drug abuse [Fillmore, 2003; Finn et al., 2000; Lyvers, 2000].

Top-Down Regulative Role of the ACC

ACC's executive role is supported by its extensive anatomical connections across different levels of the neuraxis, allowing it to integrate top-down modulatory effects within a goal-directed context. It receives sensory input, has extensive and reciprocal connections with lateral prefrontal cortex and striatum, dense projections to the motor cortex and spinal cord, and its activity is modulated by multiple neurotransmitter systems [Barbas, 2000; Devinsky et al., 1995; Paus, 2001; Picard and Strick, 2001]. Human intracranial EEG studies confirm the multiplicity of the ACC contributions. Very similar inhibitory activity has been observed in the same ACC microdomain as a function of difficulty, rarity, repetition, and errors, indicating its top-down function in regulating and implementing plans to act [Wang et al., 2005]. This finding is consistent with the proposal that ACC enhances signal relevance by exerting inhibitory influence on dlPFC, permitting goalspecific stimulus processing [Medalla and Barbas, 2009].

ACC lesions result in a variety of impairments including suppression of reflexive saccades [Paus et al., 1991], deficits in focused attention and initiation of action [Cohen et al., 1999], increased errors and impaired error correction [Alexander et al., 2007; Swick and Turken, 2002], and deficient assessment of outcomes [Kennerley et al., 2006]. By blunting the ACC activity, alcohol intoxication may partially emulate these effects, rendering a person less able to focus attention, suppress reflexive responding to irrelevant stimuli, monitor performance, or initiate purposeful behavior. In this experiment, we administered a moderate alcohol dose that people commonly consume to feel a pleasant "buzz" [Chen et al., 2004/2005]. Indeed, participants reported feeling more stimulated, high, euphoric, and sexy on the ascending BAC limb, just before performing the task. Even though alcohol intoxication blunted the ACC activity to both conflict trials and errors and resulted in increased RTs and a tendency to make errors on conflict trials, posterror slowing indicating posterror adjustment did not appear to be affected.

Impaired Cognitive Control and Impulsivity

In the present study, participants were able to marshal greater strategic control and maintain low-error rates under placebo. This was reflected in stronger ACC activity under the INC condition. ACC activity was also increased on those trials on which participants made errors, indicating proficient performance monitoring. Conversely, intoxication weakened top-down regulation on conflict trials as reflected in blunted ACC activation and a tendency to make more errors on INC condition. Alcohol also attenuated error-related ACC activity, suggesting impaired performance monitoring. In addition to a tendency to make more errors to INC stimuli, intoxication resulted in longer response times overall, concurring with other similar observations [Curtin and Fairchild, 2003; Rose and Duka, 2008]. This behavioral pattern is suggestive of a partial accuracyspeed tradeoff. Indeed, correlations between accuracy and RTs under alcohol for the two potentially conflicting conditions, CONG, and INC were -0.48 and -0.51 respectively, both P < 0.05. Posterror slowing observed in this study is consistent with other evidence [Jentzsch and Dudschig, 2009; Rabbitt, 2002]. However, there were no beverage effects on posterror RTs, suggesting that posterror adjustments are not affected by this level of acute intoxication.

Alcohol-induced tendency to make more errors has been suggested to indicate impulsive responding as alcohol impairs the ability to inhibit the prepotent responses that are no longer correct [Fillmore, 2003]. The present study provides further evidence that impulsivity may result from lowered strategic control as two indices of impulsivity (Psychoticism Scale of Eysenck's Personality Questionnaire and Impulsivity Scale on Eysenck's Impulsiveness and Venturesomeness Scale) correlated with beveragerelated differences in the number of self-corrections (trials on which participants corrected themselves). In other words, the participants with higher impulsivity scores made premature responses based on incomplete processing when intoxicated, confirming other similar reports [Marinkovic et al., 2000]. Furthermore, the number of self-corrected responses correlated with current drinking (r = 0.46, P < 0.05) and also with the Self-Rating of Effects of Alcohol reports on the number of drinks required to feel intoxicated when the person first started drinking (r = 0.53, P < 0.01), suggesting an interaction between the personality predispositions and alcohol effects. Beverage-related differences in

RTs also correlated with the current level of drinking (r = -0.60, P < 0.01), indicating that faster RTs under intoxication were related to higher regular drinking levels.

Impulsivity is a complex construct, but it is often operationalized with tasks measuring cognitive control of responses. In a series of studies using a cued go/no-go task, Fillmore and colleagues have found a dose-related increase in commission errors and slower response times to the no-go signals that were falsely preceded by a "go" cue [Marczinski and Fillmore, 2003, 2005]. Similarly, alcohol intoxication decreases cognitive control on the stopsignal task [de Wit et al., 2000; Mulvihill et al., 1997], and on a continuous performance task [Dougherty et al., 1999]. Alcohol-induced impairment is reflected in premature motor preparation based on incomplete stimulus evaluation [Marinkovic et al., 2000]. Furthermore, these effects are correlated with personality traits related to impulsivity and hyperactivity [Dougherty et al., 1999; Marinkovic et al., 2000]. In a broader sense, the underlying symptom concerns an inability to resist engaging in the activity that one declares to be unwanted or even harmful; thus, the inability to maintain inhibitory control over drinking has been considered to be fundamental to alcohol abuse [Fillmore, 2003; Finn, 2000; Jentsch and Taylor, 1999; Lyvers, 2000]. A cluster of traits termed "antisocial personality disorder," inclusive of impulsivity, hyperactivity, and sensation/novelty seeking, correlates with the early-onset of alcoholism and increased drinking [Brown et al., 1996; Finn et al., 2000; Mazas et al., 2000] as well as chronic alcohol use and dependence [Hesselbrock et al., 1985]. Vulnerability to alcoholism shares a common genetic component with antisocial personality disorder, which, as a premorbid trait, may predispose individuals to a spectrum of conduct disorders including alcohol dependence [Begleiter and Porjesz, 1999; Bowirrat and Oscar-Berman, 2005; Schuckit, 2009]. Thus, by impairing top-down regulative functions, alcohol intoxication affects cognitive evaluation of the situation and impairs finding the most suitable response strategies. It may result in disinhibited behaviors, poor-self control, and inability to desist drinking leading to a further increase in alcohol intake and tolerance. Consequently, impulsivity may mediate alcohol abuse both as a dispositional risk factor and as a consequence of excessive drinking.

Cognitive Control Is Subserved by a Distributed Network

In addition to the ACC, conflict-induced activation was observed in a distributed frontoparietal network in the present study, with the INC condition evoking stronger activity than all other conditions (Table II). The other three conditions, CONG, NEUT, and READ were differentially activated only in the left IFG. Notably, the NEUT condition evoked a stronger activity than CONG, which tended to evoke stronger activation than READ. This area's role in semantic retrieval [Wagner et al., 2001] explains its activation to NEUT (noncolor related) words, in addition to selecting among competing alternatives [Thompson-Schill et al., 1997]. Conflict-related activity was also observed in IFJ, a region which may contribute to cognitive control by setting and updating stimulus-response representations [Brass et al., 2005; Derrfuss et al., 2005]. It has also been suggested that IFJ is essential for response inhibition [Swick et al., 2008] and action observation [Binkofski and Buccino, 2006]. Its function is closely related to the intraparietal cortex (IPC), which is a part of the top-down network that is activated by allocation of attention [Corbetta, 1998]. It is thought that IPC biases processing and motor planning toward the task-relevant attribute [Wojciulik and Kanwisher, 1999] while SMA participates in motor preparation and planning especially for ambiguous tasks [Rushworth et al., 2004]. Insula has been implicated in selective attention [Corbetta et al., 1991] and working memory [Paulesu et al., 1993]. Even though all these areas are sensitive to incongruity, only the ACC was affected by alcohol intoxication. This finding suggests that this level of intoxication primarily impairs the top-down central executive centered in ACC, rather than the perceptual or motor output systems.

CONCLUSIONS AND LIMITATIONS

Better insight into the anatomical and functional properties of the ACC is essential for understanding the nature of alcohol-induced impairments, especially in the context of efforts to develop effective pharmacological treatments by targeting the relevant circuits [Myrick and Anton, 2004] and gene-neurotransmitter interactions in alcohol dependence [Bowirrat and Oscar-Berman, 2005; Schuckit, 2009]. The results of the present study indicate that moderate alcohol intoxication selectively attenuates ACC activation during high-conflict and error trials, suggesting that the regulative, top-down function is most vulnerable to the effects of alcohol intoxication. ACC is essential in guiding unrehearsed behavior, when new contingencies have to be observed and inappropriate behavior inhibited. By decreasing ACC response to conflict-inducing situations, acute intoxication decreases this self-regulatory influence on behavior, making it potentially more difficult to refrain from drinking and diminishing sound judgment. Furthermore, impulsive and erroneous responding increased under intoxication and was correlated with the current drinking levels as well as impulsivity traits. The present results support models proposing that alcohol-induced prefrontal impairments may result in diminished inhibitory control of impulsive behaviors such as alcohol abuse [Fillmore, 2003; Finn, 2000; Jentsch and Taylor, 1999; Lyvers, 2000].

In this study, effects of gender were limited to only two findings: women reported feeling more intoxicated than men, and their ACC activity was lower overall than in men. Although this observation may suggest that gender plays a minimal role in the effects of alcohol intoxication on behavioral or physiological measures, future neuroimaging studies with larger samples are needed to explore this question further [Fillmore and Weafer, 2004]. In addition, potential effects of expectancy need to be explored in future studies.

Alcohol's potentially vasoactive properties need to be considered, however, in interpreting this study's findings. The BOLD effect is expressed relative to baseline and is sensitive to local changes in cerebral blood flow (CBF), volume (CBV), and metabolic rate of oxygen (CMRO₂) [Buxton et al., 2004]. By altering hemodynamics, pharmacological agents such as alcohol may confound interpretation of the neural activation [Tracey, 2001]. Some animal studies suggest that moderately high alcohol doses do not affect CBF, CMRO₂, and glucose metabolism [Ligeti et al., 1991]. However, correct interpretation of effects of alcohol on BOLD signal requires direct measures of CBF in humans, which can be obtained noninvasively with arterial spin labeling (ASL) [Golay et al., 2004; Liu and Brown, 2007]. In a different experiment, we measured perfusion with ASL technique with the same alcohol dose as used in the present study and found minimal changes in resting CBF [Rickenbacher et al., 2010]. Nevertheless, in order to partially mitigate potential alcohol-induced baseline shifts in this study, alcohol and placebo conditions were equated at baselines by subtracting the average hemodynamic response before stimulus onset from the hemodynamic response waveforms. Future ASL studies are needed to investigate potential dose-related effects of alcohol intoxication on cerebral perfusion and BOLD contrast in order to disambiguate potential vascular confounds from neural activity [Brown et al., 2007; Liu et al., 2004].

ACKNOWLEDGMENTS

The study was carried out at Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA. We thank Sarah Sheldon for assistance.

REFERENCES

- Alexander MP, Stuss DT, Picton T, Shallice T, Gillingham S (2007): Regional frontal injuries cause distinct impairments in cognitive control. Neurology 68:1515–1523.
- APA (1994): Diagnostic and Statistical Manual of Mental Disorders DSM-IV. Washington, D.C.: American Psychiatric Association.
- Barbas H (2000): Connections underlying the synthesis of cognition, memory, and emotion in primate prefrontal cortices. Brain Res Bull 52:319–330.
- Begleiter H, Porjesz B (1999): What is inherited in the predisposition toward alcoholism? A proposed model [see comments]. Alcohol Clin Exp Res 23:1125–1135.
- Binkofski F, Buccino G (2006): The role of ventral premotor cortex in action execution and action understanding. J Physiol Paris 99:396–405.
- Bowirrat A, Oscar-Berman M (2005): Relationship between dopaminergic neurotransmission, alcoholism, and Reward Deficiency syndrome. Am J Med Genet B Neuropsychiatr Genet 132:29–37.
- Brass M, Derrfuss J, Forstmann B, von Cramon DY (2005): The role of the inferior frontal junction area in cognitive control. Trends Cogn Sci 9:314–316.

- Breslin FC, Kapur BM, Sobell MB, Cappell H (1997): Gender and alcohol dosing: A procedure for producing comparable breath alcohol curves for men and women. Alcohol Clin Exp Res 21:928–930.
- Brown GG, Perthen JE, Liu TT, Buxton RB (2007): A primer on functional magnetic resonance imaging. Neuropsychol Rev 17:107–125.
- Brown SA, Gleghorn A, Schuckit MA, Myers MG, Mott MA (1996): Conduct disorder among adolescent alcohol and drug abusers. J Stud Alcohol 57:314–324.
- Burock MA, Dale AM (2000): Estimation and detection of eventrelated fMRI signals with temporally correlated noise: A statistically efficient and unbiased approach. Hum Brain Mapp 11:249–260.
- Burock MA, Buckner RL, Woldorff MG, Rosen BR, Dale AM (1998): Randomized event-related experimental designs allow for extremely rapid presentation rates using functional MRI. Neuroreport 9:3735–3739.
- Bush G, Luu P, Posner MI (2000): Cognitive and emotional influences in anterior cingulate cortex. Trends Cogn Sci 4:215–222.
- Buxton RB, Uludag K, Dubowitz DJ, Liu TT (2004): Modeling the hemodynamic response to brain activation. Neuroimage 23 (Suppl 1):S220–S233.
- Cahalan D, Cisin IH, Crossley HM (1969): American drinking practices: A national study of drinking behavior and attitudes. Monograph #6. New Brunswick, NJ: Rutgers Center of Alcohol Studies.
- Calhoun VD, Pekar JJ, Pearlson GD (2004): Alcohol intoxication effects on simulated driving: Exploring alcohol-dose effects on brain activation using functional MRI. Neuropsychopharmacology 29:2097–2017.
- Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, Cohen JD (1998): Anterior cingulate cortex, error detection, and the online monitoring of performance. Science 280:747–749.
- Chen CM, Dufour MC, Yi H (2004/2005): Alcohol consumption among young adults ages 18–24 in the United States: Results from the 2001–2002 NESARC survey. Alcohol Res Health 28:269–280.
- Cohen J (1988): Statistical Power Analysis for the Behavioral Sciences. Hillsdale, NJ: Erlbaum.
- Cohen RA, Kaplan RF, Zuffante P, Moser DJ, Jenkins MA, Salloway S, Wilkinson H (1999): Alteration of intention and self-initiated action associated with bilateral anterior cingulotomy. J Neuropsychiatry Clin Neurosci 11:444–453.
- Coles MG, Scheffers MK, Fournier L (1995): Where did you go wrong? Errors, partial errors, and the nature of human information processing. Acta Psychol (Amst) 90:129–144.
- Corbetta M (1998): Frontoparietal cortical networks for directing attention and the eye to visual locations: Identical, independent, or overlapping neural systems? Proc Natl Acad Sci USA 95:831–838.
- Corbetta M, Miezin FM, Dobmeyer S, Shulman GL, Petersen SE (1991): Selective and divided attention during visual discriminations of shape, color, and speed: Functional anatomy by positron emission tomography. J Neurosci 11:2383–2402.
- Cox RW (1996): AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. Comp Biomed Res 29:162–173.
- Cox RW, Jesmanowicz A (1999): Real-time 3D image registration for functional MRI. Magn Reson Med 42:1014–1018.
- Curtin JJ, Fairchild BA (2003): Alcohol and cognitive control: Implications for regulation of behavior during response conflict. J Abnorm Psychol 112:424–436.

- Dale AM, Fischl B, Sereno MI (1999): Cortical surface-based analysis. I. Segmentation and surface reconstruction. Neuroimage 9:179–194.
- Dao-Castellana MH, Samson Y, Legault F, Martinot JL, Aubin HJ, Crouzel C, Feldman L, Barrucand D, Rancurel G, Feline A, Syrota A, (1998): Frontal dysfunction in neurologically normal chronic alcoholic subjects: Metabolic and neuropsychological findings. Psychol Med 28:1039–1048.
- de Wit H, Crean J, Richards JB (2000): Effects of d-amphetamine and ethanol on a measure of behavioral inhibition in humans. Behav Neurosci 114:830–837.
- Derrfuss J, Brass M, Neumann J, von Cramon DY (2005): Involvement of the inferior frontal junction in cognitive control: Metaanalyses of switching and Stroop studies. Hum Brain Mapp 25:22–34.
- Devinsky O, Morrell MJ, Vogt BA (1995): Contributions of anterior cingulate cortex to behaviour. Brain 118 (Pt 1):279–306.
- Dougherty DM, Moeller FG, Steinberg JL, Marsh DM, Hines SE, Bjork JM (1999): Alcohol increases commission error rates for a continuous performance test. Alcohol Clin Exp Res 23:1342– 1351.
- Duncan J, Owen AM (2000): Common regions of the human frontal lobe recruited by diverse cognitive demands. Trends Neurosci 23:475–483.
- Easdon C, Izenberg A, Armilio ML, Yu H, Alain C (2005): Alcohol consumption impairs stimulus- and error-related processing during a Go/No-Go Task. Brain Res Cogn Brain Res 25:873–883.
- Eysenck HJ, Eysenck SBG (1975): Manual of the Eysenck Personality Questionnaire. London: Hodder & Staughton.
- Eysenck SB, Eysenck HJ (1978): Impulsiveness and venturesomeness: Their position in a dimensional system of personality description. Psychol Rep 43(3, Pt 2):1247–1255.
- Fillmore MT (2003): Drug abuse as a problem of impaired control: Current approaches and findings. Behav Cogn Neurosci Rev 2:179–197.
- Fillmore MT, Weafer J (2004): Alcohol impairment of behavior in men and women. Addiction 99:1237–1246.
- Finn P (2000): Acute effects of alcohol on cognition and impulsive-disinhibited behavior. In: Noronha A, Eckardt M, Warren K, editors. Review of NIAAA's Neuroscience and Behavioral Research Portfolio. Bethesda, MD: US Department of Health and Human Services. pp 337–356.
- Finn PR, Sharkansky EJ, Brandt KM, Turcotte N (2000): The effects of familial risk, personality, and expectancies on alcohol use and abuse. J Abnorm Psychol 109:122–133.
- Fischl B, Sereno MI, Dale AM (1999a) Cortical surface-based analysis. II. Inflation, flattening, and a surface-based coordinate system. Neuroimage 9:195–207.
- Fischl B, Sereno MI, Tootell RB, Dale AM (1999b) High-resolution intersubject averaging and a coordinate system for the cortical surface. Hum Brain Mapp 8:272–284.
- Francis WN, Kucera H (1982): Frequency Analysis of English Usage: Lexicon and grammar. Boston: Houghton Mifflin.
- Friston KJ, Holmes AP, Price CJ, Buchel C, Worsley KJ (1999): Multisubject fMRI studies and conjunction analyses. Neuroimage 10:385–396.
- Friston KJ, Rotshtein P, Geng JJ, Sterzer P, Henson RN (2006): A critique of functional localisers. Neuroimage 30:1077–1087.
- Golay X, Hendrikse J, Lim TC (2004): Perfusion imaging using arterial spin labeling. Top Magn Reson Imaging 15:10–27.
- Gough HG (1994): Theory, development, and interpretation of the CPI socialization scale. Psychol Rep 75(1, Pt 2):651–700.

- Gundersen H, Gruner R, Specht K, Hugdahl K (2008): The effects of alcohol intoxication on neuronal activation at different levels of cognitive load. Open Neuroimag J 2:65–72.
- Hedges LV, Olkin I (1985): Statistical Methods for Meta-Analysis. Orlando: Academic Press.
- Hesselbrock VM, Hesselbrock MN, Stabenau JR (1985): Alcoholism in men patients subtyped by family history and antisocial personality. J Stud Alcohol 46:59–64.
- Holroyd CB, Coles MG (2002): The neural basis of human error processing: Reinforcement learning, dopamine, and the error-related negativity. Psychol Rev 109:679–709.
- Holroyd CB, Yeung N (2003): Alcohol and error processing. Trends Neurosci 26:402–404.
- Jentsch JD, Taylor JR (1999): Impulsivity resulting from frontostriatal dysfunction in drug abuse: Implications for the control of behavior by reward-related stimuli. Psychopharmacology (Berl) 146:373–390.
- Jentzsch I, Dudschig C (2009): Why do we slow down after an error? Mechanisms underlying the effects of posterror slowing. Q J Exp Psychol (Colchester) 62:209–218.
- Kalivas PW (2009): The glutamate homeostasis hypothesis of addiction. Nat Rev Neurosci 10:561–572.
- Kennerley SW, Walton ME, Behrens TE, Buckley MJ, Rushworth MF (2006): Optimal decision making and the anterior cingulate cortex. Nat Neurosci 9:940–947.
- Laird AR, McMillan KM, Lancaster JL, Kochunov P, Turkeltaub PE, Pardo JV, Fox PT (2005): A comparison of label-based review and ALE meta-analysis in the Stroop task. Hum Brain Mapp 25:6–21.
- Li CS, Luo X, Yan P, Bergquist K, Sinha R (2009): Altered impulse control in alcohol dependence: Neural measures of stop signal performance. Alcohol Clin Exp Res 33:740–750.
- Ligeti L, Hines K, Dora E, Sinnwell T, Huang MT, McLaughlin AC (1991): Cerebral blood flow and metabolic rate in the conscious, freely moving rat: The effects of hypercapnia, and acute ethanol administration. Alcohol Clin Exp Res 15:766–770.
- Liu TT, Brown GG (2007): Measurement of cerebral perfusion with arterial spin labeling, Part 1: Methods. J Int Neuropsychol Soc 13:517–525.
- Liu TT, Behzadi Y, Restom K, Uludag K, Lu K, Buracas GT, Dubowitz DJ, Buxton RB (2004): Caffeine alters the temporal dynamics of the visual BOLD response. Neuroimage 23:1402–1413.
- Lyvers M (2000): "Loss of control" in alcoholism and drug addiction: A neuroscientific interpretation. Exp Clin Psychopharmacol 8:225–249.
- MacLeod CM (1991): Half a century of research on the Stroop effect: An integrative review. Psychol Bull 109:163–203.
- Marczinski CA, Fillmore MT (2003): Preresponse cues reduce the impairing effects of alcohol on the execution and suppression of responses. Exp Clin Psychopharmacol 11:110–117.
- Marczinski CA, Fillmore MT (2005): Compensating for alcoholinduced impairment of control: Effects on inhibition and activation of behavior. Psychopharmacology (Berl) 181:337–346.
- Marinkovic K, Halgren E, Klopp J, Maltzman I (2000): Alcohol effects on movement-related potentials: A measure of impulsivity? J Stud Alcohol 61:24–31.
- Marinkovic K, Halgren E, Maltzman I (2004): Effects of alcohol on verbal processing: An ERP study. Alcohol Clin Exp Res 28:415–423.
- Martin CS, Earleywine M, Musty RE, Perrine MW, Swift RM (1993): Development and validation of the Biphasic Alcohol Effects Scale. Alcohol Clin Exp Res 17:140–146.

- Mazas CA, Finn PR, Steinmetz JE (2000): Decision-making biases, antisocial personality, and early-onset alcoholism. Alcohol Clin Exp Res 24:1036–1040.
- Meda SA, Calhoun VD, Astur RS, Turner BM, Ruopp K, Pearlson GD (2009): Alcohol dose effects on brain circuits during simulated driving: An fMRI study. Hum Brain Mapp 30:1257–1270.
- Medalla M, Barbas H (2009): Synapses with inhibitory neurons differentiate anterior cingulate from dorsolateral prefrontal pathways associated with cognitive control. Neuron 61:609–620.
- Mulvihill LE, Skilling TA, Vogel-Sprott M (1997): Alcohol and the ability to inhibit behavior in men and women. J Stud Alcohol 58:600–605.
- Myrick H, Anton R (2004): Recent advances in the pharmacotherapy of alcoholism. Curr Psychiatry Rep 6:332–338.
- Neumann J, Lohmann G, Derrfuss J, von Cramon DY (2005): Meta-analysis of functional imaging data using replicator dynamics. Hum Brain Mapp 25:165–173.
- Nolen-Hoeksema S, Hilt L (2006): Possible contributors to the gender differences in alcohol use and problems. J Gen Psychol 133:357–374.
- Oldfield RC (1971): The assessment and analysis of handedness: The Edinburgh inventory. Neuropsychologia 9:97–113.
- Oscar-Berman M, Marinkovic K (2007): Alcohol: Effects on neurobehavioral functions and the brain. Neuropsychol Rev 17:239–257.
- Paulesu E, Frith CD, Frackowiak RS (1993): The neural correlates of the verbal component of working memory. Nature 362:342–345.
- Paulus MP, Tapert SF, Pulido C, Schuckit MA (2006): Alcohol attenuates load-related activation during a working memory task: Relation to level of response to alcohol. Alcohol Clin Exp Res 30:1363–1371.
- Paus T (2001): Primate anterior cingulate cortex: Where motor control, drive and cognition interface. Nat Rev Neurosci 2:417–424.
- Paus T, Kalina M, Patockova L, Angerova Y, Cerny R, Mecir P, Bauer J, Krabec P (1991): Medial vs lateral frontal lobe lesions and differential impairment of central-gaze fixation maintenance in man. Brain 114 (Pt 5):2051–2067.
- Picard N, Strick PL (2001): Imaging the premotor areas. Curr Opin Neurobiol 11:663–672.
- Poldrack RA (2007): Region of interest analysis for fMRI. Soc Cogn Affect Neurosci 2:67–70.
- Procyk E, Tanaka YL, Joseph JP (2000): Anterior cingulate activity during routine and non-routine sequential behaviors in macaques. Nat Neurosci 3:502–508.
- Rabbitt P (2002): Consciousness is slower than you think. Q J Exp Psychol A 55:1081–1092.
- Raichle ME, Fiez JA, Videen TO, MacLeod AM, Pardo JV, Fox PT, Petersen SE (1994): Practice-related changes in human brain functional anatomy during nonmotor learning. Cereb Cortex 4:8–26.
- Repovs G (2004): The mode of response and the Stroop effect: A reaction time analysis. Horizons Psychol 13:105–114.
- Rickenbacher E, Greve D, Azma S, Pfeuffer K, Marinkovic K (2010): Effects of alcohol intoxication on cerebral blood flow: An Arterial Spin Labeling (ASL) Study. San Antonio, TX.
- Ridderinkhof KR, de Vlugt Y, Bramlage A, Spaan M, Elton M, Snel J, Band GP (2002): Alcohol consumption impairs detection of performance errors in mediofrontal cortex. Science 298:2209–2211.
- Ridderinkhof KR, van den Wildenberg WP, Segalowitz SJ, Carter CS (2004): Neurocognitive mechanisms of cognitive control: The role of prefrontal cortex in action selection, response inhibition, performance monitoring, and reward-based learning. Brain Cogn 56:129–140.

- Roelofs A, van Turennout M, Coles MG (2006): Anterior cingulate cortex activity can be independent of response conflict in Stroop-like tasks. Proc Natl Acad Sci USA 103:13884–13889.
- Rose AK, Duka T (2008): Effects of alcohol on inhibitory processes. Behav Pharmacol 19:284–291.
- Rushworth MF, Walton ME, Kennerley SW, Bannerman DM (2004): Action sets and decisions in the medial frontal cortex. Trends Cogn Sci 8:410–417.
- Schneider W, Chein JM (2003): Controlled and automatic processing: Behavior, theory, and biological mechanisms. Cogn Sci 27:525–559.
- Schuckit MA (2009): An overview of genetic influences in alcoholism. J Subst Abuse Treat 36:S5–S14.
- Schuckit MA, Smith TL, Tipp JE (1997): The self-rating of the effects of alcohol (SRE) form as a retrospective measure of the risk for alcoholism. Addiction 92:979–988.
- Selzer ML (1971): The Michigan alcoholism screening test: The quest for a new diagnostic instrument. Am J Psychiatry 127:1653–1658.
- Selzer ML, Vinokur A, Van Rooijen L (1975): A self-administered Short Michigan Alcoholism Screening Test (SMAST). J Stud Alcohol 36:117–126.
- Stroop J (1935): Studies of interference in serial verbal reactions. J Exp Psychol 18:643–662.
- Sullivan EV, Pfefferbaum A (2005): Neurocircuitry in alcoholism: A substrate of disruption and repair. Psychopharmacology (Berl) 180:583–594.
- Swick D, Turken AU (2002): Dissociation between conflict detection and error monitoring in the human anterior cingulate cortex. Proc Natl Acad Sci USA 99:16354–16359.
- Swick D, Ashley V, Turken AU (2008): Left inferior frontal gyrus is critical for response inhibition. BMC Neurosci 9:102.
- Tarter RE, McBride H, Buonpane N, Schneider DU (1977): Differentiation of alcoholics. Childhood history of minimal brain dysfunction, family history, and drinking pattern. Arch Gen Psychiatry 34:761–768.
- Thompson-Schill SL, D'Esposito M, Aguirre GK, Farah MJ (1997): Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation. Proc Natl Acad Sci USA 94:14792–14797.
- Tracey I (2001): Prospects for human pharmacological functional magnetic resonance imaging (phMRI). J Clin Pharmacol 41:21S–28S.
- Ullsperger M, von Cramon DY (2001): Subprocesses of performance monitoring: A dissociation of error processing and response competition revealed by event-related fMRI and ERPs. Neuroimage 14:1387–1401.
- Ullsperger M, von Cramon DY (2004): Neuroimaging of performance monitoring: Error detection and beyond. Cortex 40:593–604.
- Wagner AD, Pare-Blagoev EJ, Clark J, Poldrack RA (2001): Recovering meaning: Left prefrontal cortex guides controlled semantic retrieval. Neuron 31:329–338.
- Wang C, Ulbert I, Schomer DL, Marinkovic K, Halgren E (2005): Responses of human anterior cingulate cortex microdomains to error detection, conflict monitoring, stimulus-response mapping, familiarity, and orienting. J Neurosci 25:604–613.
- Wojciulik E, Kanwisher N (1999): The generality of parietal involvement in visual attention. Neuron 23:747–764.
- Woodward JA, Bonett DG, Brecht ML (1990): Introduction to Linear Models and Experimental Design. San Diego: Harcourt Brace Jovanovich.
- Zuckerman M (1971): Dimensions of sensation seeking. J Consult Clin Psychol 36:45–52.